

# 12TH PHYSICS

# MCQS

# ALL CHAPTERS



# **F.Sc. Physics (2<sup>nd</sup> Year) Multiple Choice Questions**

Chapter # 12: Electrostatics

Chapter # 13: Current Electricity

Chapter # 14: Electromagnetism

Chapter # 15: Electromagnetic Induction

Chapter # 16: Alternating Current

Chapter # 17: Physics of Solids

Chapter # 18: Electronics

Chapter # 19: Dawn of Modern Physics

Chapter # 20: Atomic Spectra

Chapter # 21: Nuclear Physics

## CHAPTER # 12: ELECTROSTATICS

1. Coulomb's law is only applicable for
  - a) Big charges
  - b) Small charges
  - c) Point charges
  - d) Any charges
2. The force exerted by two charged bodies on one another, obeys Coulomb's law provided that
  - a) The charges are not too small
  - b) The charges are in vacuum
  - c) The charges are not too large
  - d) The linear dimension of charges is much smaller than distance between them
3. The constant K in Coulomb's Law depends upon
  - a) Nature of medium
  - b) System of units
  - c) Intensity of charge
  - d) Both a & b
4. A unit of electric charge is:
  - a) Volt
  - b) Henry
  - c) Coulomb
  - d) Weber
5. Presence of dielectric always:
  - a) Increases the electrostatic force
  - b) Decreases the electrostatic force
  - c) Does not effect the electrostatic force
  - d) Doubles the electrostatic force
6. The S.I unit of permittivity is:
  - a)  $Nm^2/C^2$
  - b)  $C^2/Nm^2$
  - c)  $Nm/C^2$
  - d)  $Nm^2/C$
7. The lines which provide information about the electric force exerted on charged particles are:
  - a) Magnetic field lines
  - b) Electric field lines
  - c) Tangent lines
  - d) Curved lines
8. The electric field created by positive charge is:
  - a) Radially outward
  - b) Zero
  - c) Circular
  - d) Radially inward
9. The value of relative permittivity for all the dielectrics is always:
  - a) Less than unity
  - b) Greater than unity
  - c) Equal to unity
  - d) Zero
10. Photo-copier and inkjet printers are the applications of:
  - a) Electronics
  - b) Magnetism
  - c) Electrostatics
  - d) Thermodynamics
11. Selenium is a conductor material when exposed to \_\_\_\_\_
  - a) Light
  - b) Dark
  - c) Mono chromatic light
  - d) None of these
12. Selenium is an
  - a) Insulator
  - b) Conductor
  - c) Semiconductor
  - d) Photoconductor
13. In an inkjet printer, the charged ink drops are diverted by the deflection plates
  - a) Towards the charging electrodes
  - b) Towards the gutter
  - c) Towards a blank paper on which the print is to be taken
  - d) In inkjet printer ink cannot be charged
14. The electric field produced due to negative charge is always:
  - a) Radially outward
  - b) Radially inward
  - c) Circular
  - d) Zero

15. The force experience by a unit positive charge placed at a point in an electric field is called:
- Coulomb's force
  - Faraday's force
  - Lorentz's force
  - Electric field intensity
16. Of the following quantities, the one that is vector in character is an
- Electric Charge
  - Electric Field Intensity
  - Electric Energy
  - Electric Potential Difference
17. Electric field intensity is also known as
- Electric potential
  - Electric flux
  - Potential gradient
  - None
18. Potential gradient is defined as
- $\frac{\Delta E}{\Delta V}$
  - $-\frac{\Delta E}{\Delta V}$
  - $\frac{\Delta r}{\Delta V}$
  - $-\frac{\Delta V}{\Delta r}$
19. The SI unit of E are:
- (newton/meter)
  - (newton/coulomb)
  - (newton/ampere)
  - (newton  $\times$  meter)
20. The electric intensity is expressed in unit of N/C or
- Volts
  - Walt
  - Joules
  - V/m
21. The unit  $Vm^{-1}$  is equivalent to:
- $NC^{-1}$
  - $NC$
  - $NC\ m^{-1}$
  - $NmC^{-1}$
22. Electric flux is defined as:
- $\phi = \mathbf{A} \cdot \mathbf{B}$
  - $\phi = \mathbf{E} \times \mathbf{A}$
  - $\phi = \mathbf{E} \cdot \mathbf{A}$
  - $\phi = \frac{\mathbf{E}}{\mathbf{A}}$
23. When vector area is held perpendicular to the field lines, then the magnitude of electric flux is:
- Negative
  - Maximum
  - Minimum
  - Zero
24. When vector area is held parallel to electric field lines, the the magnitude of electric flux is:
- Maximum
  - Minimum
  - Either maximum or minimum
  - Negative
25. The SI unit of electric flux is:
- $NmC^{-1}$
  - $Nm^2C^{-1}$
  - $NmC^{-2}$
  - $Nm^2C^{-2}$
26. The magnitude of the electric field inside oppositely charged plates, having uniform surface charge density  $\sigma$ , is:
- $(\sigma/\epsilon_0)$
  - $(\sigma/2\epsilon_0)$
  - $(q/\epsilon_0 r)$
  - $(\sigma/2\epsilon_0 r)$
27. The electric intensity near an infinite plate of positive charge will be:
- $(q/\epsilon_0)$
  - $(\sigma/2\epsilon_0)$
  - $(q/A)$
  - $(\sigma/\epsilon_0)$
28. If a charged body is moved against the electric field, it will gain:
- Potential energy
  - Kinetic energy
  - Mechanical energy
  - None of these

29. One volt is
- One joule per coulomb
  - One dyne per coulomb
  - One Newton per coulomb
  - One watt per second
30. Absolute potential difference, due of point charge of 1C at a distance of 1m is given by:
- $9 \times 10^6 \text{ volts}$
  - $9 \times 10^7 \text{ volts}$
  - $9 \times 10^8 \text{ volts}$
  - $9 \times 10^9 \text{ volts}$
31. A charge of 0.01 C accelerated through a p.d of 1000 V acquires K.E
- 10 J
  - 100 J
  - 200 J
  - 400 eV
32. 1 joule = \_\_\_\_\_
- $6.25 \times 10^{18} \text{ eV}$
  - $6.25 \times 10^{-18} \text{ eV}$
  - $1.6 \times 10^{-19} \text{ eV}$
  - $9.1 \times 10^{-31} \text{ eV}$
33. One electron volt is equal to
- $6.25 \times 10^{18} \text{ J}$
  - $6.25 \times 10^{-18} \text{ J}$
  - $1.6 \times 10^{-19} \text{ J}$
  - $9.1 \times 10^{-31} \text{ J}$
34. How many electron will have a charge of one coulomb?
- $6.2 \times 10^{18}$
  - $6.2 \times 10^{19}$
  - $5.2 \times 10^{18}$
  - $5.2 \times 10^{19}$
35. Gravitational force between two objects does not depends on:
- Force
  - Masses
  - Distance
  - Medium
36. The charge on the electron was calculated by
- Faraday
  - J.J. Thomson
  - Millikan
  - Einstein
37. The equation for the stokes law is
- $6\pi\eta r$
  - $6\pi\eta r v$
  - $6rv$
  - $8\pi\eta r v$
38. The charge determined by the Millikan's experiment is
- $q = \frac{qv d}{m}$
  - $q = \frac{qv d}{g}$
  - $g = \frac{mg d}{v}$
  - None
39. Capacitors may be considered as a device for
- Storing energy
  - Increasing resistance
  - Decreasing resistance
  - None
40. The medium used b/w the plates of capacitor is called
- Polarization
  - Dielectric
  - Insulators
  - Medium
41. Capacity of a capacitor depends upon
- Size of plate
  - Distance b/w plates
  - Nature of dielectric b/w plates
  - All of above
42. Farad is defined as:
- $\frac{C}{V}$
  - $\frac{A}{V}$
  - $\frac{C}{J}$
  - $\frac{J}{C}$
43. The capacitance of a parallel plate capacitor is given by:
- $C = \frac{A}{\epsilon_0 d}$
  - $C = \frac{A\epsilon_0}{d}$
  - $C = \frac{\epsilon_0 d}{A}$
  - $C = \frac{d}{\epsilon_0 A}$

44. The expression of energy stored in a capacitor is given by:

- a)  $E = CV^2$
- b)  $E = \frac{1}{2} CV^2$
- c)  $E = \frac{1}{2} C^2 V$
- d)  $E = \frac{1}{2} (CV)^2$

45. Unit of energy density of electric field is:

- a)  $J C^{-1}$
- b)  $J V^{-1}$
- c)  $J m^{-3}$
- d)  $J F^{-3}$

46. The term "RC" has same unit as that of:

- a) Potential
- b) Capacitance
- c) Energy
- d) Time

47. The ratio of  $C_{vac}$  and  $C_{med}$  is equal to

- a)  $\epsilon_r$
- b)  $\frac{1}{\epsilon_r}$
- c)  $\epsilon_0$
- d)  $\frac{1}{\epsilon_0}$

48. During charging of a capacitor, the ratio of instantaneous charge and maximum charge on plates of capacitors at  $t = RC$  is

- a) 36.8%
- b) 63.2%
- c) 20%
- d) 30%

Q. No.	Ans.	Q. No.	Ans.	Q. No.	Ans.
1	c	17	c	33	c
2	d	18	d	34	a
3	d	19	b	35	d
4	c	20	d	36	c
5	b	21	a	37	b
6	b	22	c	38	c
7	b	23	d	39	a
8	a	24	a	40	b
9	b	25	b	41	d
10	c	26	a	42	a
11	a	27	b	43	b
12	d	28	a	44	b
13	b	29	a	45	c
14	b	30	d	46	d
15	d	31	a	47	b
16	b	32	a	48	b

## CHAPTER # 13: CURRENT ELECTRICITY

1. One coulomb per second is equal to
  - a) Joule
  - b) Volt
  - c) Ampere
  - d) Walt
2. In the metallic conductor the current is due to flow of \_\_\_\_\_ charge
  - a) Positive
  - b) Negative
  - c) Proton
  - d) None
3. Conventional current flow from
  - a) Point of higher potential to point of lower potential
  - b) Point of lower potential to point of higher potential
  - c) Point of lower potential to point of lower potential
  - d) None
4. In the thermocouple the heat energy is converted into
  - a) Mechanical energy
  - b) Electric energy
  - c) Magnetic energy
  - d) None
5. The heating effect of current utilized in
  - a) Iron
  - b) Tube light
  - c) Fan
  - d) Motor
6. Through an electrolyte, electric current is passed due to drift of
  - a) Free electrons
  - b) Positive and negative ions
  - c) Free electrons and holes
  - d) Protons
7. Joule law can be expressed as
  - a)  $H = I^2 R t$
  - b)  $H = I R^2 t$
  - c)  $H = \frac{I^2}{R t}$
  - d)  $H = \frac{I}{R t}$
8. The heat produced by passage of current through resistor is:
  - a)  $H = I^2 R t$
  - b)  $H = I R^2 t$
  - c)  $H = \frac{I^2}{R t}$
  - d)  $H = \frac{I}{R t}$
9. Current can be measured by using:
  - a) Heating effect
  - b) Magnetic effect
  - c) Chemical effect
  - d) None of these
10. In liquids and gases, the current is due to the motion of :
  - a) Negative charges
  - b) Positive charges
  - c) Neutral particles
  - d) Both negative and positive charges
11. When electricity passes through the liquid, then process is called:
  - a) Electro late
  - b) Electrolysis
  - c) Electro-conductor
  - d) None
12. Magnetic effect of current is utilized in
  - a) Iron
  - b) Thermocouple
  - c) Measurement of current
  - d) None
13. The VI-graph of Ohm's law is:
  - a) Hyperbola
  - b) Ellipse
  - c) Parabola
  - d) Straight
14. Mathematical form of ohm's law is
  - a)  $I = V R$
  - b)  $I = V / R$
  - c)  $I = R / V$
  - d)  $R = I V$
15. Ohm's law is valid for only current flowing in
  - a) Conductors
  - b) Transistors
  - c) Diodes
  - d) Electric Areas

16. The proportionality constant between current and potential difference is:
- $\rho$
  - $R$
  - $C$
  - $V$
17. 1 ohm is defined as:
- $\frac{V}{C}$
  - $\frac{V}{A}$
  - $\frac{A}{C}$
  - $\frac{V}{A}$
18. In series circuit the net resistance is
- Algebraic Sum of all resistance
  - Sum of reciprocals of all resistances in circuit
  - Remain constant
  - None
19. The reciprocal of resistivity is called
- Resistance
  - Conduction
  - Conductivity
  - None
20. The unit of conductivity is
- $\Omega \cdot m$
  - $(\Omega \cdot m)^{-1}$
  - $\Omega \cdot m^{-1}$
  - None
21. A wire of resistance  $R$  is cut into two equal parts, its resistance becomes  $R/2$ . What happens to resistivity?
- Double
  - Same
  - Half
  - One forth
22. When temperature increases, the resistance of conductor:
- Increases
  - Decreases
  - Remains constant
  - Vanishes
23. Heat sensitive resistors are called
- Resistors
  - Capacitors
  - Thermistors
  - Inductors
24. Thermistor can be used for the accurate measurement of
- Voltage
  - Resistance
  - Temperature
  - Heat
25. The maximum power delivered by battery is:
- $P_{max} = \frac{E^2}{4r}$
  - $P_{max} = 4rE^2$
  - $P_{max} = VIT$
  - Unlimited
26. If the length and diameter of conductor is double, the resistance is
- Remain same
  - Double
  - Half
  - Four times
27. A wire of uniform cross-section  $A$  and length  $L$  is cut into two equal parts. The resistance of each part becomes:
- Double
  - Half
  - 4 times
  - $\frac{1}{4}$  times
28. The fractional change in resistivity per Kelvin
- Co-efficient in resistance
  - Co-efficient of resistivity
  - Resistance
  - None
29. In the carbon resistor their value can be find by their
- Wires
  - Terminals
  - Color Bands
  - Spots
30. The third band is written in the form of power of
- 2
  - 6
  - 8
  - 10
31. The numerical value of black color is:
- 3
  - 2
  - 1
  - 0

32. The color code for the color Grey is  
a) 7  
b) 8  
c) 9  
d) 5
33. The colors of strips on a certain carbon resistor from extreme left are yellow, black and red respectively. Its resistance is:  
a) 4 k $\Omega$   
b) 400  $\Omega$   
c) 40  $\Omega$   
d) 40 k $\Omega$
34. If the tolerance color is gold then its value is  
a)  $\pm 2\%$   
b)  $\pm 4\%$   
c)  $\pm 5\%$   
d)  $\pm 6\%$
35. Tolerance for silver band is:  
a)  $\pm 5\%$   
b)  $\pm 10\%$   
c)  $\pm 15\%$   
d)  $\pm 20\%$
36. A rheostat can be used as a  
a) Variable resistor  
b) Potential divider  
c) Both a and b  
d) None of these
37. The wire used in Rheostat is made from  
a) Constantan  
b) Nichrome  
c) Manganin  
d) Tungsten
38. The S.I unit of emf is same as:  
a) Work  
b) Energy  
c) Power  
d) Potential Difference
39. The terminal potential difference of a battery of internal resistance "r" and emf " $\varepsilon$ " is:  
a)  $V = \varepsilon + Ir$   
b)  $V = \varepsilon - Ir$   
c)  $V = \frac{\varepsilon - r}{I}$   
d)  $V = \frac{I}{\varepsilon - r}$
40. Which electric bulb has the least resistance?  
a) 60 watts  
b) 100 watts  
c) 200 watts  
d) 500 watts
41. An electric heater 220V, 440W has a resistance  
a) 2  $\Omega$   
b) 110  $\Omega$   
c) 0.5  $\Omega$   
d) 20  $\Omega$
42. Kirchhoff's first rule is:  
a)  $\sum V = 0$   
b)  $\sum R = 0$   
c)  $\sum I = 0$   
d)  $\sum T = 0$
43. Kirchhoff's first rule is based on conservation of:  
a) Energy  
b) Voltage  
c) Charge  
d) Mass
44. The algebraic sum of all the current at junction is zero, is Kirchhoff's  
a) 1<sup>st</sup> law  
b) 2<sup>nd</sup> law  
c) 3<sup>rd</sup> law  
d) 4<sup>th</sup> law
45. The algebraic sum of voltages changes around a closed circuit or loop is zero, is Kirchhoff's  
a) 1<sup>st</sup> law  
b) 2<sup>nd</sup> law  
c) 3<sup>rd</sup> law  
d) 4<sup>th</sup> law
46. An ideal voltmeter would have an infinite  
a) Current  
b) Voltage  
c) Resistance  
d) None of these
47. The emf of two cells can be compared by  
a) AVO meter  
b) Voltmeter  
c) Potentiometer  
d) Galvanometer

48. An accurate measurement of emf of a cell is made by
- A voltmeter
  - An ammeter
  - A potentiometer
  - All of them

49. The ratio of emf of two cells  $\varepsilon_1/\varepsilon_2$ , is equal to
- $l_1/l_2$
  - 1 : 2
  - $l_2/l_1$
  - 2 : 1

Q. No.	Ans.	Q. No.	Ans.	Q. No.	Ans
1	c	18	a	35	b
2	b	19	c	36	c
3	a	20	b	37	c
4	b	21	b	38	d
5	a	22	a	39	b
6	b	23	c	40	d
7	a	24	c	41	b
8	a	25	a	42	c
9	b	26	c	43	c
10	d	27	b	44	a
11	b	28	b	45	b
12	c	29	c	46	c
13	d	30	d	47	c
14	b	31	d	48	c
15	a	32	c	49	a
16	b	33	d		
17	d	34	c		

## CHAPTER # 14. ELECTROMAGNETISM

- The units of magnetic field B, in system international is:
  - Weber
  - Tesla
  - Gauss
  - Newton
- One tesla (T) is:
  - $1T = 1N A m^{-1}$
  - $1T = 1N A^{-1} m^{-1}$
  - $1T = 1N A m$
  - $1T = 1N m A^{-1}$
- The magnetic flux " $\Phi$ " through an area " $A$ " is:
  - $\Phi = \mathbf{B} \times \mathbf{A}$
  - $\Phi = \mathbf{B} \cdot \mathbf{A}$
  - $\Phi = \mathbf{A} \times \mathbf{B}$
  - None of these
- One Tesla is also equal to
  - $\text{wb.m}^2$
  - $\text{wb.m}^{-2}$
  - $\text{wb.m}$
  - None
- Torque on a current carrying coil is:
  - $BINA \cos \alpha$
  - $BINA \sin \alpha$
  - $BIL \cos \alpha$
  - $BIL \sin \alpha$
- The magnetic force is simply a:
  - Reflecting force
  - Deflecting force
  - Restoring force
  - Gravitational force
- The galvanometer can be made sensitive if the value of the factor  $\frac{C}{BNA}$  is:
  - Made large
  - Made small
  - Remains constant
  - Infinite
- When a small resistance is connected parallel to the galvanometer, the resulting circuit behaves as:
  - Voltmeter
  - Wheatstone bridge
  - Ammeter
  - Potentiometer
- The anode in the CRO is:
  - Control number of electrons
  - Control the brightness of spot formed
  - Accelerates and focus the beam
  - At negative potential with respect to cathode
- The galvanometer constant in a moving coil galvanometer is given by:
  - $K = \frac{NB}{CA}$
  - $K = \frac{C}{NAB}$
  - $K = \frac{NAB}{C}$
  - $K = \frac{CA}{NB}$
- $\sum_{r=1}^N \mathbf{B} \cdot \mathbf{dL} = \mu_0 I$  is the relation for:
  - Milikan's law
  - Gauss's law
  - Ampere's law
  - Lenz's law
- The brightness of spot on CRO screen is controlled by:
  - Anodes
  - Cathodes
  - Grid
  - Plates
- To measure the current in a circuit, ammeter is always connected in:
  - Parallel
  - Series
  - Sometimes parallel sometimes series
  - Neither series nor parallel
- If the angle b/w  $\vec{v}$  and B is zero then magnetic force will be
  - Max
  - Min
  - Zero
  - None

15. A charged particles is projected at an angle into a uniform magnetic field. Which of the following parameter of the charged particle will be affected by magnetic field:
- Energy
  - Momentum
  - Speed
  - Velocity
16. Force on a moving charge in a uniform magnetic field will be maximum, when angle between  $v$  and  $B$  is:
- $0^\circ$
  - $30^\circ$
  - $60^\circ$
  - $90^\circ$
17. The S.I. unit of magnetic flux is
- Tesla
  - Weber
  - Joule
  - Newton
18. Beam of electrons are also called:
- Positive rays
  - x-rays
  - cathode rays
  - cosmic rays
19. Tesla is the unit of
- Electric field
  - Magnetic field
  - Magnetic field intensity
  - Electric field intensity
20. It is possible to set a charge at rest into motion with magnetic field
- Yes
  - No
  - Some Time
  - None
21. The grid in CRO \_\_\_\_\_
- Controls the number of electrons accelerated by anode
  - Controls the brightness of the spot fall on the screen
  - Both a and b
  - Deflects the beam of electrons
22. To convert a Weston-type galvanometer into voltmeter, the series resistance is given by \_\_\_\_\_
- $R_h = \frac{V}{I_g}$
  - $R_h = \frac{V}{I_g} - R_g$
  - $R_h = \frac{V}{R_g} - I_g$
  - None of these
23. The shape of magnetic field around a long straight current carrying wire is
- Electrical
  - Squire
  - Varies with current
  - Circular
24. The electrons of mass " $m$ " and charge " $e$ " is moving in a circle of radius " $r$ " with speed " $v$ " in a uniform magnetic field of strength " $B$ ". then
- $r \propto m$
  - $r \propto B$
  - $r \propto \frac{1}{v}$
  - $r \propto \frac{1}{m}$
25. The toque in the coil can be increased by increasing
- Number of turns
  - Current and magnetic field
  - Area of coil
  - All of above
26. A current carrying loop, when placed in a uniform magnetic field will experience
- Electric flux
  - Torque
  - Magnetic flux
  - Force
27. The magnetic flux will be maximum if the angle between magnetic field strength and vector area is:
- $0^\circ$
  - $60^\circ$
  - $90^\circ$
  - $180^\circ$
28. One weber is equal to
- $N.A^2/A$
  - $N.m^2/A$
  - $N.A/m$
  - $N.m/A$

29. The waveform of sinusoidal voltage, its frequency and phase can be found by  
a) CRO  
b) Diode  
c) Transistor  
d) Radio
30. The force on a charge particle moving parallel to magnetic field is  
a) Maximum  
b) Minimum  
c) Zero  
d) None
31. The unit of permeability of free space is  
a) T.m/A  
b) T.m<sup>2</sup>/A  
c) T.m/A<sup>2</sup>  
d) None
32. The value of  $\mu_o$  is  
a)  $4 \pi \times 10^{-6}$   
b)  $4 \pi \times 10^{-7}$   
c)  $4 \pi \times 10^{-8}$   
d)  $4 \pi \times 10^{-9}$
33. The magnetic induction inside current carrying solenoid is  
a)  $\mu_o nI$   
b)  $\mu_o NL$   
c)  $\mu_o N$   
d) None
34.  $F = F_e + F_m$  is  
a) Electric force  
b) Magnetic force  
c) Lorentz force  
d) None
35. The material used in fluorescent screen is  
a) Electric  
b) Magnetic  
c) Phosphors  
d) None
36. In the galvanometer the current is proportional to  
a) Magnetic field  
b) Electric field  
c) Angle  
d) None
37. When a small resistance is connected in parallel to the galvanometer it is called  
a) Ammeter  
b) Voltmeter  
c) AVO meter  
d) None
38. The relation between current " $I$ " and deflection " $\theta$ " in a moving coil galvanometer is:  
a)  $I \propto \frac{1}{\theta}$   
b)  $I \propto \cos \theta$   
c)  $I \propto \sin \theta$   
d)  $I \propto \theta$
39. To convert a galvanometer into voltmeter we connect a resistance in  
a) Series  
b) Parallel  
c) Series or parallel  
d) None
40. AVO-meter is used to find  
a) Current  
b) Voltage  
c) Resistance  
d) All of above
41. An ideal voltmeter has  
a) Small resistance  
b) High resistance  
c) Infinite resistance  
d) None
42. A galvanometer can be more sensitive if C/BAN is made  
a) Very large  
b) Very small  
c) Unaltered  
d) None
43. Ammeter and galvanometer  
a) Are always connected in series  
b) Are always connected in parallel  
c) Both in series and parallel  
d) None

44. The sensitivity of galvanometer is directly depends on  
 a) Magnetic field  
 b) Area of coil  
 c) Number of turns  
 d) All of above
45. The dot product of magnetic field induction and vector area is called  
 a) Electric flux  
 b) Magnetic flux  
 c) Ampere law  
 d) None
46. When the number of turns in a solenoid is doubled without any change in the length of the solenoid its self induction will be:  
 a) Four times  
 b) Doubled  
 c) Halved  
 d) None
47. The wave form of sinusoidal voltage, its frequency and phase can be found by  
 a) CRO  
 b) Diode  
 c) Transistor  
 d) Radio
48. Voltmeter is used to measure:  
 a) Current  
 b) Resistance  
 c) Temperature  
 d) Potential difference
49. The resistance of a voltmeter should have a very high resistance  
 a) It does not disturb the circuit  
 b) It draws some current  
 c) It same the galvanometer coil  
 d) None of these
50. A voltmeter is always connected in:  
 a) Parallel  
 b) Series  
 c) Perpendicular  
 d) Straight line

51. To find the shunt resistance we used equation

$$\begin{aligned} \text{a) } R_s &= \frac{I_g R_g}{I - I_g} \\ \text{b) } R_s &= \frac{I_s R_g}{I - I_g} \\ \text{c) } R_s &= \frac{I_g R_s}{R - I_g} \\ \text{d) } R_s &= \frac{I_s R_s}{I - I_g} \end{aligned}$$

52. Ammeter is used to measure:

- a) Resistance  
 b) Voltage  
 c) Current  
 d) Capacitance

53. An avo-meter is also called:

- a) An ammeter  
 b) A voltmeter  
 c) A multi.meter  
 d) An ohm-meter

### Key Chapter # 14

Q. No.	Ans.	Q. No.	Ans.	Q. No.	Ans.
1	b	19	b & c	37	a
2	b	20	b	38	d
3	b	21	c	39	a
4	b	22	b	40	d
5	a	23	d	41	c
6	b	24	a	42	b
7	b	25	d	43	a
8	c	26	b	44	d
9	c	27	a	45	b
10	b	28	d	46	a
11	c	29	a	47	a
12	c	30	c	48	d
13	b	31	a	49	a
14	c	32	b	50	a
15	b & d	33	a	51	a
16	d	34	c	52	c
17	b	35	c	53	c
18	c	36	c		

## CHAPTER # 15. ELECTROMAGNETIC INDUCTION

1. The induced e.m.f. is produced due to
  - a) Motion of coil
  - b) Motion of magnet
  - c) The rate of change of flux
  - d) None
2. The direction of induced current is always so as to oppose the change which causes the current is called:
  - a) Faraday's law
  - b) Lenz's law
  - c) Ohm's law
  - d) Kirchhoff's 1<sup>st</sup> rule
3. The energy stored per unit volume inside a solenoid is calculated by:
  - a)  $\frac{1}{2} \frac{B^2}{\mu_0} (Al)$
  - b)  $\frac{1}{2} \frac{B^2}{\mu_0}$
  - c)  $\frac{1}{2} \frac{\mu_0}{B^2} (Al)$
  - d)  $\frac{1}{2} \frac{\mu_0}{B^2}$
4. The SI units of induced emf is
  - a) Ohm
  - b) Tesla
  - c) Henry
  - d) Volt
5. The principle of an alternating current generator is based on:
  - a) Coulomb's law
  - b) Ampere's law
  - c) Faraday's law
  - d) Lenz's law
6. If velocity of a conductor moving through a magnetic field B is made zero, then motional emf is:
  - a)  $-vBL$
  - b)  $-\frac{v}{BL}$
  - c)  $-\frac{BL}{v}$
  - d) Zero
7. If we make the magnetic field stronger, the value of induced current is:
  - a) Decreased
  - b) Increased
  - c) Vanished
  - d) Kept constant
8. The inductance is more in self induction in:
  - a) Air cored coil
  - b) Iron cored coil
  - c) Tungsten cored coil
  - d) None of these
9. One henry is equal to:
  - a)  $1 \text{ ohm} \times 1 \text{ second}$
  - b)  $1 \text{ ohm} \times 1 \text{ meter}$
  - c)  $1 \text{ ohm} \times 1 \text{ coulomb}$
  - d) None of these
10. A device which converts electrical energy into mechanical energy is called:
  - a) Transformer
  - b) AC generator
  - c) DC motor
  - d) DC generator
11. When constant current flows in primary of transformer, then the emf induced across secondary of transformer is:
  - a) Zero
  - b) Constant
  - c) Alternating
  - d) Irregular
12. 1 henry
  - a)  $1 \text{ V A s}^{-1}$
  - b)  $1 \text{ V s A}^{-1}$
  - c)  $1 \text{ V m A}^{-1}$
  - d)  $1 \text{ V A m}^{-1}$
13. A generator converts mechanical energy into
  - a) Chemical energy
  - b) Light energy
  - c) Heat energy
  - d) Electrical energy
14. When a loop of wire is moved across a magnetic field, the current is produced in it is called
  - a) Eddy current
  - b) Direct current
  - c) Photo electric current
  - d) Induced current

15. Energy stored in an inductor is:
- $\frac{1}{2}LI^2$
  - $\frac{1}{2}LI$
  - $\frac{1}{2}L^2I^2$
  - $\frac{1}{2}LI$
16. If fingers of right hand show the direction of magnetic field and palm shows the direction of force, then thumb points for:
- Torque
  - Voltage
  - Current
  - Induced emf
17. Induced electric current can be explained using which law
- Gauss's law
  - Faraday's law
  - Ohm's law
  - Ampere law
18. Lenz's law is consistent with law of conservation of
- Mass
  - Energy
  - Charge
  - None
19. An inductor is a circuit element that can store energy in the form of
- Magnetic field
  - Electric flux
  - Electric field
  - None
20. The negative sign with induced e.m.f. is due to
- Faraday's law
  - Lenz's law
  - Ampere law
  - None
21. The relation of motional e.m.f. , when a conductor is move in perpendicular magnetic field, is:
- $E=BLV$
  - $E=qBl$
  - $E=Blq$
  - $E=qVB$
22. If we increase the resistance of the circuit containing a coil, the induced e.m.f. will be
- Increase
  - Decrease
  - Remain same
  - None
23. The self-inductance may be defined by
- $L = \frac{-\epsilon}{\Delta I / \Delta t}$
  - $L = \frac{-\Delta I / \Delta t}{\epsilon}$
  - $L = \frac{-\epsilon}{\Delta \phi / \Delta t}$
  - $L = \frac{\epsilon}{\Delta \phi / \Delta t}$
24. Inductance are measured by
- Coulombs
  - Amperes
  - Volt
  - Henry
25. An over loaded motor draws
- Max. current
  - Min. current
  - Half
  - None
26. The co-efficient of mutual inductance is equal to
- $\epsilon \left( \frac{\Delta I_p}{\Delta t} \right)$
  - $\epsilon \left( \frac{\Delta t}{\Delta I_p} \right)$
  - $\epsilon \Delta I_p \Delta t$
  - None
27. Alternating current changes
- Its magnitude as well as direction
  - Only direction but not magnitude
  - Only magnitude but not direction
  - None
28. Inductance is measured in:
- Volt
  - Ampere
  - Henry
  - Ohm

29. The instantaneous value of A.C. voltage is
- $V = V_0 \sin 2\pi ft$
  - $V = V_0 \sin 2\pi t$
  - $V = V_0 \sin 2\pi wt$
  - None
30. The induced e.m.f. in A.C. generator is
- $VBL \sin \phi$
  - $NBSN \sin \phi$
  - $NAB \sin \phi$
  - $NIAB \sin \phi$
31. The back motor effect exist in the
- Generator
  - Mater
  - A.C. Meter
  - None
32. The coil used in the generators is called
- Commutaters
  - Slip rings
  - Armature
  - None
33. The back ward generator is called
- Electric motor
  - A.C. generator
  - Reverse generator
  - None
34. The principle of transformer is
- Amperes law
  - Mutual induction
  - Motional e.m.f.
  - None
35. A transformer is a device which step up or stop down
- Energy
  - Power
  - Voltage
  - All of above
36. An ideal transformer obeys the law of conservation of:
- Flux
  - Momentum
  - Emf
  - Energy
37. The coil which is connected to input of a transformer is called:
- Primary
  - Secondary
  - Middle
  - None
38. In the actual transformer, the output is always
- Equal to input
  - Less then input
  - More than input
  - None
39. In ideal transformer when applied potential difference is double, the current is:
- Doubled
  - Tripled
  - Halved
  - Same
40. For a good transformer the hysteries loop are \_\_\_\_\_ in size.
- Small
  - Large
  - Zero
  - None
41. To minimize the heating effect in the transmission lines
- High current, low voltage in used
  - High voltage, low current in used
  - Same voltage and current in used
  - None
42. Maximum emf generated in a generator is:
- $\varepsilon = \varepsilon_0 \sin \theta$
  - $\varepsilon = N\omega AB \sin \theta$
  - $\varepsilon = N\omega AB$
  - None of these
43. Induced e.m.f is
- Directly proportional to change in flux
  - Directly proportional to rate of change of flux
  - Inversely proportional to change of flux
  - None of these

44. Lenz's law is in accordance with the law of conservation of:
- Momentum
  - Angular momentum
  - Energy
  - Charge
45. When motor is at its Max. speed the back e.m.f will be
- Maximum
  - Zero
  - Cannot tell
  - None of these
46. The application of mutual induction is a
- Television
  - Radio
  - D.C. motor
  - Transformer
47. The ratio of average induced emf to the rate of change of current in the coil is called:
- Self inductance
  - Mutual inductance
  - Self inductance
  - Mutual inductanc
48. Which of the following is not present in AC generator:
- Armature
  - Magnet
  - Slip rings
  - Commutator

## Key Chapter # 15

Q. No.	Ans.	Q. No.	Ans.	Q. No.	Ans.
1	c	17	b	33	a
2	b	18	b	34	b
3	a	19	a	35	c
4	d	20	b	36	d
5	c	21	a	37	a
6	d	22	c	38	b
7	b	23	a	39	c
8	b	24	d	40	a
9	a	25	a	41	b
10	c	26	b	42	c
11	a	27	a	43	b
12	b	28	c	44	c
13	d	29	a	45	a
14	d	30	none	46	d
15	a	31	a	47	a
16	c	32	c	48	d

## CHAPTER # 16. ALTERNATING CURRENT CIRCUITS

1. The mean value of A.C. over a complete cycle in
  - a) Maximum
  - b) Minimum
  - c) Zero
  - d) None
2. The inductive reactance is:
  - a)  $X_L = \omega C$
  - b)  $X_L = \omega L$
  - c)  $X_L = \frac{1}{\omega C}$
  - d)  $X_L = \frac{1}{\omega L}$
3. At high frequency, the current through a capacitor of AC circuit will be:
  - a) Large
  - b) Small
  - c) Infinite
  - d) Zero
4. The highest value reached by the voltage or current in one cycle is called
  - a) Peak to peak value
  - b) Peak value
  - c) Instantaneous value
  - d) Root mean square value
5. If the motor is overloaded, then the magnitude of "back emf":
  - a) Increases
  - b) Decreases
  - c) Constant
  - d) Becomes zero
6. A capacitor is perfect insulator for:
  - a) Alternating current
  - b) Direct current
  - c) Both a and b
  - d) None
7. The process combining low frequency signal with high frequency radio wave is called:
  - a) Modulation
  - b) Amplification
  - c) Demodulation
  - d) Resonance
8. In pure resistive AC circuit, instantaneous value of voltage or current:
  - a) Current lags behind voltage
  - b) Current leads voltage by  $\frac{\pi}{2}$
  - c) Both are in-phase
  - d) Voltage leads current by  $\frac{\pi}{2}$
9. Which of the following requires a material medium for their propagation:
  - a) Heat waves
  - b) X-rays
  - c) Sound waves
  - d) Ultraviolet rays
10. In modulation, low frequency signal is known as:
  - a) Loaded signal
  - b) Fluctuated signal
  - c) Harmonic signal
  - d) Modulation signal
11. The mutual induction between two coils depends upon:
  - a) Area of the coils
  - b) Number of turns
  - c) Distance between the coils
  - d) All of these
12. Pure choke consumes:
  - a) Minimum power
  - b) Maximum power
  - c) No power
  - d) Average power
13. To construct a step down transformer:
  - a)  $N_S < N_P$
  - b)  $N_P < N_S$
  - c)  $N_S = N_P$
  - d)  $N_S \cdot N_P = 1$
14. Power dissipation in pure inductive or in a pure capacitive circuit is:
  - a) Infinite
  - b) Zero
  - c) Minimum
  - d) Maximum
15. The practical application of phenomenon of mutual induction is:
  - a) Electrical motor
  - b) Transformer
  - c) AC generator
  - d) DC generator
16. There are \_\_\_\_\_ types of modulations:
  - a) 1
  - b) 2
  - c) 3
  - d) 4

17. The SI unit of impedance is:
  - a) Henry
  - b) Hertz
  - c) Ampere
  - d) Ohm
18. Which of the following is true for a step down transformer:
  - a)  $N_p > N_s$
  - b)  $N_p < N_s$
  - c)  $N_p = N_s$
  - d) *None of these*
19. If  $I_0$  is the peak value of AC, its average value over the complete cycle is:
  - a)  $\frac{I_0}{\sqrt{2}}$
  - b)  $\sqrt{2}I_0$
  - c)  $\frac{\sqrt{2}}{I_0}$
  - d) *zero*
20. At resonance, the phase angle for RLC series resonance circuit equals:
  - a)  $0^\circ$
  - b)  $90^\circ$
  - c)  $180^\circ$
  - d)  $270^\circ$
21. The unit of impedance is:
  - a) *Ohm*
  - b)  $(Ohm)^{-1}$
  - c)  $(Ohm - m)^{-1}$
  - d) *no unit*
22. A device that allows permits flow of DC through the circuit easily, is called:
  - a) Inductor
  - b) Capacitor
  - c) AC generator
  - d) Transformer
23. The phase difference between each pair of coils of a three phase AC generator is:
  - a)  $0^\circ$
  - b)  $90^\circ$
  - c)  $120^\circ$
  - d)  $180^\circ$
24. Main reason for the world wide use of AC is that it can be transmitted to:
  - a) Short distances at very low cost
  - b) Long distances at very high cost
  - c) Short distances at very high cost
  - d) Long distances at very low cost
25. If  $V_{rms} = 10\sqrt{2}$  volts, then phase voltage  $V_0$  will be:
  - a) *10 volts*
  - b) *20 volts*
  - c) *40 volts*
  - d)  $\frac{10}{\sqrt{2}}$  volts
26. At resonance RLC series circuit shows the behavior of:
  - a) Pure resistive circuit
  - b) Pure capacitive circuit
  - c) Pure inductive circuit
  - d) Pure RLC circuit
27. At resonance, the value of current in RLC series circuit is equal to:
  - a)  $\frac{V_0}{R}$
  - b)  $V_0 R$
  - c)  $\frac{I}{2}$
  - d) *zero*
28. At high frequency, RLC series circuit shows the behavior of:
  - a) Pure inductive circuit
  - b) Pure resistive circuit
  - c) Pure capacitive circuit
  - d) Pure RLC circuit
29. The r.m.s. value of A.C current in
  - a)  $0.707 I_0$
  - b)  $0.707 V_0$
  - c)  $0.707 R_0$
  - d) *None*
30. In pure resistive A.C. circuit the voltage and current are
  - a) *In phase*
  - b) Voltage leads the current
  - c) Current leads the voltage
  - d) *None*
31. The waves which can also pass through the vacuum are
  - a) Matter wave
  - b) Mechanical wave
  - c) Electromagnetic wave
  - d) Transverse wave
32. The unit used for capacitive reactance is
  - a) Volt
  - b) Ampere
  - c) Joule
  - d) Ohm

33. Power dissipated in pure inductor is:  
 a) Large  
 b) Small  
 c) Infinite  
 d) Zero
34. If the frequency of A.C in large the reactance of capacitor is  
 a) Large  
 b) Small  
 c) Zero  
 d) None
35. In case of capacitor, the voltage lag behind the current by  
 a)  $90^\circ$   
 b)  $60^\circ$   
 c)  $30^\circ$   
 d)  $180^\circ$
36. In the pure inductor the resistance is  
 a) Zero  
 b) Maximum  
 c) Minimum  
 d) None
37. In pure inductive circuit the voltage  
 a) Lead the current by  $90^\circ$   
 b) Ledge the current by  $90^\circ$   
 c) Remain same with current  
 d) None
38. The reactance of inductor is represented by  
 a)  $X_C$   
 b)  $X_L$   
 c)  $R_L$   
 d) None
39. If the frequency of A.C. is doubled, the reactance of inductor will be  
 a) Half  
 b) Same  
 c) Double  
 d) Triple
40. The average power dissipated in a pure inductor is  
 a) Maximum  
 b) Minimum  
 c) Zero  
 d) None
41. By increasing the frequency of A.C. through an inductor the reactance will be  
 a) Increases  
 b) Decreases  
 c) Remain same  
 d) None
42. In case of phasor diagram the vector rotates  
 a) Clockwise  
 b) Anti clockwise  
 c) Remain stationary  
 d) None
43. The combine opposition of resistor, capacitor and inductor is called  
 a) Reactance  
 b) Resistor  
 c) Impedance  
 d) None
44. The S.I unit of impedance is called  
 a) Joule  
 b) Weber  
 c) Ampere  
 d) Ohm
45. When A.C. flow through RC series circuit the magnitude of voltage is  
 a)  $V = I\sqrt{R^2 + X_L^2}$   
 b)  $V = I\sqrt{R^2 + X_C^2}$   
 c)  $V = IR$   
 d) None
46. The magnitude of voltage in case of RL – series circuit  
 a)  $V = I\sqrt{R^2 + X_L^2}$   
 b)  $V = I\sqrt{R^2 + X_C^2}$   
 c)  $V = IR$   
 d) None
47. The average power in case of A.C. series circuit is  
 a)  $P = VI$   
 b)  $P = VI \cos \phi$   
 c)  $P = VI \sin \phi$   
 d) None
48. In equation  $P = VI \cos \phi$ , the factor  $\cos \phi$  is called  
 a) Cosine factor  
 b) Power factor  
 c) Phase  
 d) None
49. The behavior of resistance is frequency  
 a) Dependent  
 b) Independent  
 c) No, response  
 d) None of these

50. The impedance  $Z$  can be expressed as:
- $V_{rms} + I_{rms}$
  - $V_{rms} - I_{rms}$
  - $V_{rms} / I_{rms}$
  - $I_{rms} / V_{rms}$
51. At resonance frequency the power factor is
- One
  - Zero
  - Two
  - Three
52. The frequency at which  $X_L$  is equal to  $X_C$  is called
- Resonance frequency
  - Threshold frequency
  - Non-frequency
  - None
53. At resonance frequency the impedance of A.C series circuit is
- Maximum
  - Minimum
  - Can not explain by give data
  - None
54. In parallel RLC circuit, at resonance frequency, there will be maximum
- Power
  - Voltage
  - Impedance
  - None
55. The electrical oscillators are used in
- Metal detectors
  - Amplifier
  - Diode
  - None
56. Which of the following permits direct current to flow easily?
- Resistance
  - Capacitance
  - Inductance
  - None of these
57. A.M stands for
- Amplitude Modulation
  - Applied Metal
  - Accurate Measurement
  - None
58. F.M stands for
- Frequency Modulation
  - Frequency Metal
  - Frequency Member
  - None
59. The process of combining the low frequency signal with high frequency radio-wave is called
- Modulation
  - Amplification
  - Rectification
  - None
60. A capacitor is perfect insulator for:
- Alternating current
  - Direct current
  - Both a and b
  - None
61. During each cycle A.C voltage reaches its peak value
- One time
  - Two times
  - Four times
  - None of these
62. In modulation, high frequency radio wave is called:
- Fluctuated wave
  - Carrier wave
  - Matter wave
  - Energetic wave
63. At high frequency the reactance of the capacitor is
- Low
  - Large
  - Very large
  - None of these
64. The behavior of resistance is frequency
- Dependent
  - Independent
  - No response
  - None of these
65. In an inductor the phase difference between the current and voltage is
- Current lags voltage by  $90^\circ$
  - Voltage lags current by  $180^\circ$
  - Current leads voltage by  $90^\circ$
  - None of these
66. The condition of resonance reached when
- $X_C > X_L$
  - $X_L < X_C$
  - $X_L = X_C$
  - None of these

67. The phase difference between coils of three phase A.C is  
 a)  $60^\circ$   
 b)  $45^\circ$   
 c)  $90^\circ$   
 d)  $120^\circ$
68. Modulation is the process in which  
 a) Amplitude is change  
 b) Frequency is change  
 c) Both a & b  
 d) None of these
69. How many times per second will an incandescent lamp reach maximum brilliance when connected to a 50Hz source?  
 a) 50 times  
 b) 100 times  
 c) 200 times  
 d) None of these
70. The peak value of sinusoidal voltage in an AC circuit is 50V. The rms value of voltage is roughly equal to  
 a) 70V  
 b) 40V  
 c) 35V  
 d) 45V
71. In RLC series AC circuit, when  $X_L = X_C$  then impedance is  
 a) Minimum  
 b) Maximum  
 c) Zero  
 d) None

## Key Chapter # 16

Q. No.	Ans.	Q. No.	Ans.	Q. No.	Ans.
1	c	25	b	49	b
2	b	26	a	50	c
3	a	27	a	51	a
4	b	28	c	52	a
5	b	29	a	53	b
6	b	30	a	54	c
7	a	31	c	55	a
8	c	32	d	56	c
9	c	33	d	57	a
10	d	34	b	58	a
11		35	a	59	a
12	c	36	a	60	b
13	a	37	a	61	b
14	b	38	b	62	b
15	b	39	c	63	a
16	b	40	c	64	b
17	d	41	a	65	a
18	a	42	b	66	c
19	d	43	c	67	d
20	a	44	d	68	c
21	a	45	b	69	b
22	a	46	a	70	c
23	c	47	b	71	a
24	d	48	b		

## CHAPTER # 17. PHYSICS OF SOLIDS

1. What is the S.I unit of modules of elasticity of substances?
  - a)  $\text{Nm}^{-2}$
  - b)  $\text{Jm}^{-2}$
  - c)  $\text{Nm}^{-1}$
  - d) Being number, it has no unit
2. The bands in atom containing conductive electrons, according to “band theory of solids” is
  - a) Conduction band
  - b) Valance band
  - c) Forbidden band
  - d) None of these
3. The substances which have partially filled conduction bands are called:
  - a) Insulators
  - b) Semi-conductors
  - c) Conductors
  - d) Super conductors
4. What type of impurity is to be added to the semi-conductor material to provide holes:
  - a) Monovalent
  - b) Trivalent
  - c) Tetravalent
  - d) Pentavalent
5. Which of the following is an example of ductile substances:
  - a) Lead
  - b) Copper
  - c) Glass
  - d) Lead and copper
6. When a stress changes the shape of a body, it is called:
  - a) Volumetric stress
  - b) Shear stress
  - c) Tensile stress
  - d) Compressional strees
7. The ration of applied stress to volumetric strain is called:
  - a) Young's modulus
  - b) Shear modulus
  - c) Bulk modulus
  - d) Tensile modulus
8. Substances which break just after the elastic limit is reached are called:
  - a) Ductile substances
  - b) Hard substances
  - c) Soft substances
  - d) Brittle substances
9. Cure temperature for iron is:
  - a)  $0^\circ\text{C}$
  - b)  $570^\circ\text{C}$
  - c)  $750^\circ\text{C}$
  - d)  $1025^\circ\text{C}$
10. The SI unit of stress is same as that of:
  - a) Momentum
  - b) Pressure
  - c) Force
  - d) Length
11. Which of the following has least energy gap?
  - a) Conductors
  - b) Insulators
  - c) Semi-conductors
  - d) None of these
12. A magnetism produced by electrons within an atom is due to:
  - a) Spin motion of electrons
  - b) Orbital motion of electrons
  - c) Both Spin and orbital motion of electrons
  - d) Vibratory motion of electrons
13. If the conductivity of a material is high, then it is:
  - a) An insulator
  - b) A semi-conductor
  - c) A good conductor
  - d) A super condutor
14. A substance having empty conduction band is called:
  - a) Semi-conductor
  - b) Conductor
  - c) Insulator
  - d) None of these
15. The stress that produces change in length is known as:
  - a) Tensile stress
  - b) Shear stress
  - c) Volumetic stress
  - d) Longitudenal stress

16. What are the dimensions of stress?  
a)  $MLT^{-2}$   
b)  $ML^{-2}T^{-1}$   
c)  $ML^{-1}T^{-2}$   
d)  $ML^0T^{-1}$
17. Which one of the following physical quantities does not have the dimensions of force per unit?  
a) Stress  
b) Strains  
c) Young's modulus  
d) Pressure
18. Germanium is:  
a) semi-conductor  
b) conductor  
c) insulator  
d) none of these
19. unit of strain is:  
a)  $\frac{N}{m^2}$   
b)  $\frac{N}{m}$   
c)  $Nm$   
d) *no unit*
20. At curie temperature, iron becomes:  
a) Ferromagnet  
b) Diamagnet  
c) Paramagnet  
d) Super-conductor
21. Materials that undergo plastic deformation before breaking are called \_\_\_\_\_  
a) Brittle  
b) Ductile  
c) Amorphous  
d) Polymers
22. Formation of large molecule by joining small molecules is \_\_\_\_\_  
a) Fusion  
b) Polymerization  
c) Crystallization  
d) Subtraction
23. Any alteration produced in shapes, length or volume when a body is subjected to some external force is called \_\_\_\_\_  
a) Stiffness  
b) Ductility  
c) extension  
d) deformation
24. The energy band occupied by the valence electrons is called \_\_\_\_\_  
a) Energy state  
b) Valence band  
c) -ve energy state  
d) Conduction band
25. the substances having negative temperature coefficient of resistance is called:  
a) Conductors  
b) Insulators  
c) Semi-conductor  
d) None of these
26. The Curie temperature is that at which \_\_\_\_\_  
a) Semi conductor becomes conductors  
b) Ferromagnetic becomes paramagnetic  
c) Paramagnetic becomes diamagnetic  
d) Metal becomes super conductor
27. Materials in which valence electrons are tightly bound to their atoms at low temperature are called \_\_\_\_\_  
a) Semi conductors  
b) Super conductors  
c) Insulators  
d) Conductors
28. The band theory of solids explains satisfactorily the nature of  
a) Electrical insulators alone  
b) Electrical conductors alone  
c) Electrical semi conductors alone  
d) All of the above
29. A vacant or partially filled band is called \_\_\_\_\_  
a) Conduction band  
b) Valence band  
c) Forbidden band  
d) Empty band
30. A completely filled or partially filled band is called \_\_\_\_\_  
a) Conduction band  
b) Valence band  
c) Forbidden band  
d) Core band

31. Which one has the greatest energy gap  
\_\_\_\_\_
- Semi-conductors
  - Conductors
  - Metals
  - Non-metals
32. With increase in temperature, the electrical conductivity of intrinsic semi conductors  
\_\_\_\_\_
- Decreases
  - Increases
  - Remain the same
  - First increases, then decreases
33. Holes can exists in \_\_\_\_\_
- Conductors
  - Insulators
  - Semi conductors
  - All of the above
34. In a semi conductors, the charge carriers are \_\_\_\_\_
- Holes only
  - Electrons only
  - Electrons and holes both
  - All of the above
35. The net charge on N-type material is  
\_\_\_\_\_
- Positive
  - Negative
  - Both a & b
  - Neutral
36. The most stable material for making permanent magnet is:
- Iron
  - Steel
  - Aluminum
  - Copper
37. Pentavalent impurities are called
- Donor impurities
  - Acceptor impurities
  - Sometimes donor and some times
  - Acceptors
38. Minority carriers in N-type materials are
- Electrons
  - Protons
  - Neutrons
  - Holes
39. The temperature at which conductors lose its resistivity is called
- Supper temperature
  - Kelvin temperature
  - Critical temperature
  - None
40. The magnetic domains are the small regions of the order of
- Millimeter
  - Micrometer
  - Micron
  - None of these
41. N-type semi-conductor is obtained by doping intrinsic semi-conductors with \_\_\_\_\_
- Tetravalent impurity atom
  - Trivalent impurity atom
  - Pentavalent impurity atom
  - Hexavalent impurity atom
42. The first supper conductor was discovered by
- Fermi
  - Kmaerling
  - Weinberg
  - None
43. Examples of brittle substances are
- Glass
  - Copper
  - Lead
  - None
44. Example of crystalline solids are also
- Metals
  - Ionic compounds
  - Ceramics
  - All of them
45. a semi-conductor will behave as insulator when:
- High potential difference is applied
  - When its temperature is 0 k
  - Pentavalent impurity added
  - Trivalent impurity added
46. The field of long bar magnet is like a
- Solenoid
  - Toroid
  - Pieces of magnet
  - None

47. The curie temperature of Iron is  
 a) 600°C  
 b) 650°C  
 c) 700°C  
 d) 750°C
48. The examples of diamagnetic are  
 a) Water  
 b) Copper  
 c) Antimony  
 d) All of them
49. Strain is dimensionless and has  
 a) Units  
 b) No units  
 c) S.I units  
 d) None
50. The electrons occupying the outermost shell of an atom and the electrons occupying in the energy band are called  
 a) Energy band  
 b) Valence band  
 c) Forbidden energy band  
 d) None of these
51. Conductors are those materials in which the free electrons  
 a) Very large  
 b) Very small  
 c) Plenty of  
 d) None of these
52. The magnetism produced by electrons within an atom is due to  
 a) Spin motion  
 b) Orbital motion  
 c) Spin & orbital motion  
 d) None of these
53. The combination of solenoid and a specimen of iron inside it make a powerful magnet called  
 a) Horse shoe magnet  
 b) Bar magnet  
 c) Electromagnet  
 d) None of these
54. the substance in which the atoms don't form magnetic dipole are called:  
 a) ferromagnetic  
 b) paramagnetic  
 c) diamagnetic  
 d) conductors
55. A current which demagnetize the material completely is called  
 a) Applied current  
 b) Coercive current  
 c) Maximum current  
 d) None of these
56. The energy need to magnetize and demagnetize the specimen during the each cycle of magnetizing current is  
 a) Value of current  
 b) Value of demagnetizing current  
 c) Value of magnetic flux density  
 d) Area of the loop
57. The temperature below which resistivity of some materials becomes zero, is called:  
 a) Kelvin temperature  
 b) Critical temperature  
 c) Absolute zero temperature  
 d) Limiting temperature
58. A well known example of an intrinsic semi-conductor is:  
 a) Germanium  
 b) Phosphorous  
 c) Aluminum  
 d) Cobalt
59. The critical temperature for mercury is:  
 a) 7.2 K  
 b) 4.2 K  
 c) 1.18 K  
 d) 3.7 K

## Key Chapter # 17

Q. No.	Ans.	Q. No.	Ans.	Q. No.	Ans.
1	a	21	b	41	c
2	a	22	b	42	b
3	c	23	d	43	a
4	c	24	b	44	d
5	b	25	c	45	a
6	b	26	b	46	a
7	c	27	a	47	d
8	d	28	d	48	d
9	c	29	a	49	b
10	b	30	b	50	b
11	a	31	d	51	a
12	c	32	b	52	c
13	c	33	c	53	c
14	c	34	c	54	c
15	a	35	d	55	b
16	c	36	b	56	d
17	b	37	a	57	b
18	a	38	d	58	a
19	d	39	c	59	b
20	c	40	a	60	c

## CHAPTER # 18. ELECTRONICS

1. A semi – conductor can be used as a rectifier because
  - a) It has low resistance to the current flow when forward biased
  - b) It has high resistance to the current flow when reversed biased
  - c) It has low resistance to the current flow when forward biased and high resistance when reversed biased
  - d) None of the above
2. The central region of a transistor is called:
  - a) Base
  - b) Emitter
  - c) Collector
  - d) Neutral
3. The SI unit of current gain is:
  - a) Ampere
  - b) Volt
  - c) Ohm-meter
  - d) It has no units
4. A NAND gate with two inputs A & B has an output 0 if
  - a) A is 0
  - b) B is 0
  - c) Both A and B are 0
  - d) Both A and B are 1
5. The gain G of non inverting operational amplifier is
  - a)  $G = -\frac{R_2}{R_1}$
  - b)  $G = 1 + \frac{R_2}{R_1}$
  - c)  $G = \frac{R_2}{R_1}$
  - d)  $G = 1 - \frac{R_2}{R_1}$
6. When a PN junction is reverse biased, the depletion region is
  - a) Widened
  - b) Narrowed
  - c) Normal
  - d) No change
7. A potential barrier of 0.7 V exist across pn junction made from
  - a) Silicon
  - b) Germanium
  - c) Indium
  - d) Gallium
8. A diode characteristics curve is a graph plotted between
  - a) Current and time
  - b) Voltage and time
  - c) Voltage and current
  - d) Forward voltage and reverse current
9. The output of AND gate will be 1 when
  - a) Both inputs are at 0
  - b) Either one input is at 1
  - c) Both inputs are at 1
  - d) None of these
10. For non-inverting amplifiers if  $R_1 = \infty \Omega$  and  $R_2 = 0 \Omega$ , then gain of amplifier is
  - a) -1
  - b) 0
  - c) +1
  - d) infinite
11. An expression for current gain of a transistor is given by
  - a)  $\beta = \frac{I_B}{I_C}$
  - b)  $\beta = I_B + I_C$
  - c)  $\beta = I_C - I_B$
  - d)  $\beta = \frac{I_C}{I_B}$
12. In n-type materials, the minority carriers are
  - a) Free electrons
  - b) Holes
  - c) Protons
  - d) Mesons
13. Transistors are made from
  - a) Plastics
  - b) Metals
  - c) Insulators
  - d) Doped semi-conductors
14. The number of diodes in a bridge rectifier is
  - a) 4
  - b) 2
  - c) 3
  - d) 5
15. The reverse current through PN junction is
  - a) Infinite
  - b) Zero
  - c) Less than forward current
  - d) Greater than forward current
16. Photocells are used for
  - a) Security system
  - b) Counting system
  - c) Automatic door system
  - d) All of these

17. Transistor has  
a) 2 regions  
b) 3 regions  
c) 4 regions  
d) 1 region
18. A complete amplifier circuit made on a silicon chip and enclosed in a small capsule is called  
a) Diode  
b) Inductor  
c) Resistor  
d) Operational amplifier
19. The open loop gain of an operational amplifier is of the order of  
a)  $10^8$   
b)  $10^5$   
c)  $10^2$   
d)  $10^{-3}$
20. The automatic working of streets lights is due to  
a) Inductor  
b) Capacitor  
c) Comparator  
d) Rectifier
21. In half wave rectification, the output DC voltage is obtained across the load for  
a) The positive half cycle of input AC  
b) The negative half cycle of input AC  
c) The positive and negative half cycles of input AC  
d) Either positive or negative half cycle of input AC
22. The color of light emitted by LED depends on  
a) Its forward biased  
b) Its reversed biased  
c) The amount of forward current  
d) The type of semi conductor material used
23. A PN junction photodiode is  
a) Operated in forward direction  
b) Operated in reversed direction  
c) A very fast photo detector  
d) Dependent on thermally generated minority carriers
24. The reverse current through semi-conductor diode is due to  
a) Holes  
b) Electrons  
c) Majority carriers  
d) Minority carriers
25. The potential barrier for PN junction made from Si at room temperature is  
a) 0.9 V  
b) 0.3 V  
c) 0.7 V  
d) 0.8 V
26. Process of conversion of DC to AC is called  
a) Rectification  
b) Amplification  
c) Oscillation  
d) Modulation
27. Which one of the following is called fundamental gate  
a) NOR gate  
b) NOT gate  
c) NAND gate  
d) Exclusive OR gate
28. For proper working of a transistor in normal circuits  
a) Emitter base junction is reversed biased, collector base junction is forward biased  
b) Emitter base junction is forward biased and collector base junction is forward biased  
c) C-B junction is reversed biased, E-B junction is forward biased  
d) C-B junction is reversed biased and E-B junction is reversed biased
29. In a properly biased NPN transistor most of the electrons from the emitter  
a) Recombine with holes in the base  
b) Recombine in the emitter itself  
c) Pass through the base to the collector  
d) Are stopped by the junction barrio
30. A diode characteristic curve is a graph between  
a) Current and time  
b) Voltage and time  
c) Voltage and current  
d) Forward voltage and reverse current
31. A NOR Gate is ON only when all its input are  
a) ON  
b) OFF  
c) Positive  
d) High
32. A logic gate is an electronic circuit which  
a) Makes logic decision  
b) Work on binary algebra  
c) Alternates between 0 and 1  
d) None of these
33. The output of a 2-input OR gate is zero only when its  
a) Both input are zero  
b) Either input is 1  
c) Both input are 1  
d) Either input is 0

34. An XOR gate produces an positive logic output only when its two inputs are  
a) High  
b) Low  
c) Different  
d) Same
35. An AND Gate  
a) Implement logic addition  
b) Is equivalent to a series switching circuit  
c) Is any or all gate  
d) Is equivalent to a parallel switching circuit
36. The only function of a NOT gate is to  
a) Stop a signal  
b) Re-complement a signal  
c) Invert an input signal  
d) Acts as a universal gate
37. The forward current through a semiconductor diode circuit is due to  
a) Minority carriers  
b) Majority carriers  
c) Holes  
d) Electrons
38. The device used for conversion of AC into DC is  
a) An oscillator  
b) A detector  
c) An amplifier  
d) A rectifier
39. The thickness of depletion region is of the order of  
a)  $10^{-7}$  m  
b)  $10^{-6}$  m  
c)  $10^{-5}$  m  
d)  $10^{-4}$  m
40. The ratio of  $\beta$  gives the  
a) Voltage gain  
b) Current gain  
c) Input resistance  
d) None
41. The resistance between + ive and – ive inputs of op – amplifier is  
a)  $100 \Omega$   
b)  $1000 \Omega$   
c)  $10^6 \Omega$   
d) None of these
42. Photo – voltaic cell have  
a) Battery input  
b) No external bias  
c) No internal bias  
d) None
43. Transistor can be used as  
a) Oscillators  
b) Switches  
c) Memory unit  
d) All of them
44. NOT gate has only  
a) One input  
b) Two inputs  
c) Many inputs  
d) None
45. A photo – diode can switch its current ON and OFF in  
a) Milli seconds  
b) Micro seconds  
c) Nano seconds  
d) None
46. Diode is a device which has \_\_\_\_\_ terminals.  
a) One  
b) Two  
c) Three  
d) Four
47. Transistor is a device which has \_\_\_\_\_ terminals.  
a) One  
b) Two  
c) Three  
d) Four
48. The Boolean expression  $X = A + B$  represents the logic operation of  
a) NAND gate  
b) NOR gate  
c) OR gate  
d) NOT gate
49. The open loop gain of op – amplifier is  
a) Zero  
b) High  
c) Very high  
d) Low
50. The width of depletion region of a diode  
a) Increases under forward bias  
b) Is independent of applied voltage  
c) Increases under reverse bias  
d) None of these
51. A LED emits lights only  
a) Forward biased  
b) Reverse Biased  
c) Un biased  
d) None of these

52. NAND gate is a combination
- AND gate and NOT gate
  - AND gate and OR gate
  - OR gate and NOT gate
  - NOT gate and NOT gate
53. The reverse or leakage current of the diode is of the order of
- Microampere
  - Milli-ampere
  - Both
  - None of these
54. Temperature, pressure etc are converted into electronic informations by devices called
- LEDs
  - Sensors
  - Vacuum tubes
  - None
55. Base of the transistor is very thin of the order of
- $10^{-2}m$
  - $10^{-4}m$
  - $10^{-6}m$
  - $10^{-8}m$
56. How many diodes are used for the full wave bridge rectifier circuit is
- Two
  - Three
  - Four
  - None of these
57. The electronic circuits which implement the various logic operations are known as
- Digital gates
  - Logic gate
  - Voltage operated gate
  - All of them
58. In a half-wave rectifier the diode conducts during
- Both halves of the input cycle
  - A portion of the positive half of the input cycle
  - A portion of the negative half of the input cycle
  - One half of the input cycle
59. The output of a two inputs OR gate is 0 only when its
- Both inputs are 0
  - Either input is 1
  - Both inputs are 1
  - Either input is zero

60. For typical transistor as an amplifier

$$a) \frac{V_{out}}{V_{in}} = \beta \frac{R_C}{R_{ie}}$$

$$b) \frac{V_{out}}{V_{in}} = \beta$$

$$c) \frac{V_{out}}{V_{in}} = \beta \frac{R_{ic}}{R_c}$$

$$d) \frac{V_{out}}{V_{in}} = \beta \frac{R_{ie}}{R_{ie}}$$

61. The resistance between (+) and (-) of ideal Op-Amp is
- High
  - Low
  - Infinity
  - Moderate

### Key Chapter # 18

Q. No.	Ans.	Q. No.	Ans.	Q. No.	Ans.
1	c	22	d	43	d
2	a	23	b	44	a
3	d	24	d	45	c
4	d	25	c	46	b
5	b	26	a	47	c
6	a	27	b	48	c
7	a	28	c	49	c
8	c	29	c	50	c
9	c	30	c	51	a
10	c	31	b	52	a
11	d	32	a	53	a
12	b	33	a	54	b
13	d	34	c	55	c
14	a	35	b	56	c
15	c	36	c	57	b
16	d	37	b	58	d
17	b	38	d	59	a
18	d	39		60	a
19	b	40	b	61	a
20	c	41	c		
21	d	42	b		

## CHAPTER # 19. DAWN OF MODERN PHYSICS

1. The Einstein mass-energy relationship is
  - a)  $E = mc$
  - b)  $E = mc^3$
  - c)  $E = mc^2$
  - d)  $E = m^2c^2$
2. When an electron combines with a positron, we get
  - a) One photon
  - b) Two photons
  - c) Three photons
  - d) Four photons
3. Production of X-rays can be regarded as the reverse phenomenon of
  - a) Pair production
  - b) Photoelectric effect
  - c) Compton effect
  - d) Annihilation of matter
4. The radius of atom is the order of
  - a)  $10^{10} m$
  - b)  $10^{-10} m$
  - c)  $10^{-14} m$
  - d)  $10^{14} m$
5. In 1905, the special theory of relativity was proposed by
  - a) Maxwell
  - b) De Broglie
  - c) Bohr
  - d) Einstein
6. Neutron was discovered in 1932 by
  - a) Bohr
  - b) Chadwick
  - c) Dirac
  - d) Fermi
7. The rest mass of photon is
  - a) Infinity
  - b) Zero
  - c)  $hf$
  - d)  $mc^2$
8. A maximum Compton shift in the wavelength of scattered photon will occur at
  - a)  $\theta = 0^\circ$
  - b)  $\theta = 45^\circ$
  - c)  $\theta = 90^\circ$
  - d)  $\theta = 180^\circ$
9. The Davisson and Germer experiment indicates
  - a) Interference
  - b) Polarization
  - c) Electron diffraction
  - d) Refraction
10. A positron is a particle having
  - a) Mass equal to electron
  - b) Charge equal to electron
  - c) Equal mass but opposite charge to electron
  - d) Mass equal to proton
11. In Compton scattering, the Compton shift  $\Delta\lambda$  will be equal to Compton wavelength if the scattering angle is
  - a)  $0^\circ$
  - b)  $45^\circ$
  - c)  $60^\circ$
  - d)  $90^\circ$
12. Unit of Planck's constant is
  - a) volt
  - b) J s
  - c)  $J s^{-1}$
  - d) eV
13. Which one is most energetic?
  - a)  $\gamma$  - rays
  - b) X - rays
  - c) Ultraviolet rays
  - d) Visible light
14. The total amount of energy radiated per unit orifice area of cavity radiator per unit time is directly proportional to
  - a)  $T$
  - b)  $T^2$
  - c)  $T^3$
  - d)  $T^4$
15. Planck's constant  $h$  has the same units as that of
  - a) Linear momentum
  - b) Angular momentum
  - c) Torque
  - d) Power
16. Photoelectric effect was explained by
  - a) Hertz
  - b) Einstein
  - c) Rutherford
  - d) Bohr
17. All motions are
  - a) Absolute
  - b) Uniform
  - c) Relative
  - d) Variable
18. The rest mass energy of an electron in MeV is equal to
  - a) 0.511
  - b) 0.611
  - c) 0.902
  - d) 1.02

19. An observer shoots parallel to a meter stick at very high speed (relativistic) and finds that the length of meter stick is \_\_\_\_\_  
a) Greater than one meter  
b) Less than one meter  
c) One meter  
d) None of these
20. Linear momentum of a photon is  
a) Zero  
b)  $hf/c^2$   
c)  $hf/c$   
d)  $c^2/hf$
21. Photon with energy greater than 1.02 MeV can interact with matter as  
a) Photoelectric effect  
b) Compton effect  
c) Pair production  
d) Pair annihilation
22. Stopping potential for a metal surface in case of photo electric emission depends on  
a) The threshold frequency for the metal surface  
b) The intensity of incident light  
c) The frequency of incident light and the work function for metal surface  
d) None of these
23. As the temperature of black body is raised, the wavelength corresponding to maximum intensity  
a) Shifts towards longer wavelength  
b) Shifts towards shorter wavelength  
c) Remains the same  
d) Shifts towards shorter as well as longer wavelength
24. The name of photon for quantum of light was proposed by  
a) Ampere  
b) Planck's  
c) Thomson  
d) Einstein
25. A photon is a \_\_\_\_\_  
a) Unit of energy  
b) Positively charged particle  
c) Packet of electromagnetic radiations  
d) Unit of wavelength
26. The light of suitable frequency falling on metal surface ejects electrons, this phenomenon is called  
a) X-ray emission  
b) Compton effect  
c) Photoelectric effect  
d) Nuclear fission
27. The minimum energy needed for a photon to create an electron-positron pair is  
a) 1.02 KeV  
b) 0.51 KeV  
c) 0.51 MeV  
d) 1.02 MeV
28. Davisson and Germer indicates \_\_\_\_\_ in their experiment  
a) Electron refraction  
b) Electron polarization  
c) Electron reflection  
d) Electron diffraction
29. In Davison – Germer experiment, the diffracted proton from crystal shows \_\_\_\_\_  
a) Particle property  
b) Wave property  
c) Light property  
d) Quantum property
30. In electron microscope, electric and magnetic field are used as \_\_\_\_\_  
a) Electromagnetic gun  
b) Source of electromagnetic waves  
c) Deflected charged particle  
d) Converging source of electrons
31. The uncertainty in momentum and position is due to its \_\_\_\_\_  
a) Property of matter and radiation  
b) Two dimensional motions  
c) Motion of certain wave length  
d) Very high velocity
32. The energy radiated is directly proportional to fourth power of Kelvin's temperature is \_\_\_\_\_  
a) Karl-wein's laws  
b) Raleigh jeans law  
c) Stephens law  
d) Planck's
33. The anti-particle of electron is  
a) Proton  
b) Positron  
c) Meson  
d) Neutron
34. The reverse process of pair-production is  
a) Annihilation  
b) Materialization  
c) Fission  
d) Fusion
35. The decrease in length with speed was explained by  
a) Einstein  
b) Lorentz  
c) Bohr  
d) None

36. All the motion in this universe are  
 a) Absolute  
 b) Uniform  
 c) Variable  
 d) Relative
37. Pair production cannot possible in  
 a) Air  
 b) Water  
 c) Glass  
 d) Vacuum
38. The minimum energy required for pair production is  
 a) 10.2 Mev  
 b) 1.02 Mev  
 c) 102 Mev  
 d) None
39. The relation  $\lambda_{Max}T = Contt.$  is  
 a) Wein's Law  
 b) Plank's Law  
 c) Stephen Law  
 d) None
40. A quantity  $\sqrt{1 - \frac{v^2}{c^2}}$  is always  
 a) Greater than one  
 b) Less than one  
 c) Equal to one  
 d) None of these
41. Who gave the idea of matter wave?  
 a) De-Broglie  
 b) Planck  
 c) Einstein  
 d) Huygen
42. The Stefan-Boltzmann's constant has the value  
 a)  $5.67 \times 10^{-5} \text{Wm}^{-2}\text{K}^{-4}$   
 b)  $5.67 \times 10^{-6} \text{Wm}^{-1}\text{K}^{-4}$   
 c)  $5.67 \times 10^{-6} \text{Wm}^{-2}\text{K}^{-4}$   
 d)  $5.67 \times 10^{-8} \text{Wm}^{-2}\text{K}^{-4}$
43. The energy of photon of radio waves is only about  
 a)  $10^{-6} \text{eV}$   
 b)  $10^{-4} \text{eV}$   
 c)  $10^{-10} \text{eV}$   
 d)  $10^{-12} \text{eV}$
44. The idea of quantization of energy was proposed by  
 a) Einstein  
 b) Max Planck  
 c) Compton  
 d) None of these
45. Application of photoelectric effect is  
 a) Photo diode  
 b) Photo transistor  
 c) Photocell  
 d) None of these
46. In Compton effect, the law/laws are conserved  
 a) Energy  
 b) Momentum  
 c) Both  
 d) None of these
47. The equations of pair production is  
 a)  $hf = 2m_0c^2 - KE(e^-) + K.E(e^+)$   
 b)  $hf = 2m_0c^2 + KE(e^-) + K.E(e^+)$   
 c)  $hf = 2m_0^2c^2 + KE(e^-) + K.E(e^+)$   
 d)  $hf = 2m_0^2c + KE(e^-) + K.E(e^+)$
48. Which of the following has the same dimension as  $h/m_0c$ ?  
 a) Length  
 b) Time  
 c) Mass  
 d) None
49. Photon 'A' has twice the energy of photon 'B'. What is the ratio of the momentum of 'A' to that of 'B'?  
 a) 4 : 1  
 b) 2 : 1  
 c) 1 : 2  
 d) None
50. Electron is an antiparticle of  
 a) Proton  
 b) Photon  
 c) Positron  
 d) Deuteron

## Key Chapter # 19

Q. No.	Ans.	Q. No.	Ans.	Q. No.	Ans.
1	c	18	a	35	a
2	b	19	b	36	d
3	b	20	c	37	d
4	b	21	c	38	b
5	d	22	a	39	a
6	b	23	b	40	b
7	b	24	d	41	a
8	d	25	c	42	d
9	c	26	c	43	c
10	c	27	d	44	b
11	d	28	d	45	c
12	b	29	b	46	c
13	a	30	d	47	b
14	d	31	a	48	a
15	b	32	c	49	b
16	b	33	b	50	c
17	c	34	a		

## CHAPTER # 20: ATOMIC SPECTRA

- Which is an example of continuous spectra?
  - Black body radiation
  - Molecular spectra
  - Atomic spectra
  - None of these
- Line spectra is an example of
  - Atomic
  - Molecular
  - Black body radiation
  - None of these
- The unit of Rydberg's constant  $R_H$  is:
  - $m^{-2}$
  - $m^{-1}$
  - $m^1$
  - $m^2$
- In a meta-stable state an can reside for about:
  - $10^{-8}s$
  - $10^{-10}s$
  - $10^{-9}s$
  - $10^{-3}s$
- Which of the following series of H-spectrum lies in ultraviolet region:
  - Lyman series
  - Balmer series
  - Paschen series
  - Bracket series
- The reverse process of photoelectric effect is:
  - Compton effect
  - X-rays production
  - Pair production
  - Pair annihilation
- Helium-Neon laser discharge tube contains Neon equal to:
  - 25%
  - 40%
  - 15%
  - 82%
- The value of Rydberg constant is:
  - $1.0974 \times 10^7 m^{-1}$
  - $1.0794 \times 10^7 m^{-1}$
  - $1.0974 \times 10^9 m^{-1}$
  - $1.974 \times 10^7 m^{-1}$
- The relation between Rhdberg constant  $R_H$  and ground state energy  $E_0$  is given by:
  - $R_H = \frac{E_0}{hc}$
  - $R_H = \frac{hc}{E_0}$
  - $E_0 = \frac{R_H}{hc}$
  - $R_H = E_0 hc$
- The radius of 3<sup>rd</sup> Bohr orbit in H-atom is greater than the radius of 1<sup>st</sup> orbit by the factor
  - 2
  - 3
  - 4
  - 9
- The orbital angular momentum in the allowed stationary orbits of H-atom is given by:
  - $\frac{2\pi}{nh}$
  - $\frac{nh}{2\pi}$
  - $\frac{2\pi}{2h}$
  - $\frac{n\pi}{h}$
- If one or more electrons are completely removed from an atom then the atom is said to be:
  - Excited
  - Polarized
  - Stablized
  - Ionized
- The quantized radius of first bohr orbit of H-atom is:
  - 0.053 nm
  - 0.0053 nm
  - 0.00053 nm
  - 53 nm
- When an electron absorbs energy, it jumps to:
  - Lower energy state
  - Higher energy state
  - Ground energy state
  - Remains in the same state
- LASER light has the property of:
  - Coherent waves
  - Non-coherent waves
  - Sound waves
  - Water waves

16. Excited atoms return to their ground state in \_\_\_\_\_  
a)  $10^{-10}$  s  
b)  $10^{-8}$  s  
c)  $10^{-6}$  s  
d)  $10^{-9}$  s
17. X-rays are \_\_\_\_\_  
a) Unknown nature  
b) High energy electrons  
c) High energy photon  
d) Radioisotopes
18. Total number of series in hydrogen spectrum is \_\_\_\_\_  
a) Three  
b) Four  
c) Five  
d) Six
19. The radiations emitted from hydrogen filled discharge tube show \_\_\_\_\_  
a) Bound spectrum  
b) Line spectrum  
c) Continuous spectrum  
d) Absorption spectrum
20. If the ionization energy of H-atom is 13.6 eV, its ionization potential will be:  
a) 13.6 V  
b) 136.0 V  
c) 3.4 V  
d) None of these
21. Radiation with wavelength longer than red light \_\_\_\_\_  
a) Ultraviolet rays  
b) X-rays  
c) Infrared radiation  
d) Visible radiations
22. Bracket series is obtained when all transition of electron terminate on \_\_\_\_\_  
a) 4<sup>th</sup> orbit  
b) 5<sup>th</sup> orbit  
c) 3<sup>rd</sup> orbit  
d) 2<sup>nd</sup> orbit
23. X - rays are similar in nature to \_\_\_\_\_  
a) Cathode rays  
b) Positive rays  
c)  $\gamma$  - rays  
d)  $\alpha$  - rays
24. The characteristic X-rays spectrum is due to  
a) The illumination of the target metal by ultraviolet radiation  
b) The bombardment of the target by proton  
c) The bombardment of target by electron  
d) The absorption of Y-radiation by the target metal
25. Wave like characteristic of electron is demonstrated by \_\_\_\_\_  
a) Line spectrum of atoms  
b) Production of X-rays  
c) Diffraction by crystalline solids  
d) Photo electric effect
26. In laser production, the state in which more atoms are in the upper state than in the lower one is called \_\_\_\_\_  
a) Metal stable state  
b) Normal state  
c) Inverted population  
d) All the above
27. Reflecting mirrors in laser is used to \_\_\_\_\_  
a) Further stimulation  
b) Lasing more  
c) For production more energetic laser  
d) All the above
28. The velocity of laser light is \_\_\_\_\_  
a) Less than ordinary light  
b) More than ordinary light  
c) Equal to ordinary light  
d) Different for different colors or frequency
29. X – rays is also known as  
a) Photon  
b)  $\gamma$  – rays  
c) Breking radiation  
d) none
30. Which one of the following is more coherent  
a) X – rays  
b) Normal light  
c) Laser  
d)  $\gamma$  – rays
31. Sunlight spectrum is  
a) Discrete  
b) Line spectrum  
c) Continuous spectrum  
d) None

32. Optical pumping exist in  
a) X – rays  
b) Laser  
c) Spectrum  
d) None
33. The total energy of electron in an orbit around the nucleus is  
a) + ive  
b) ive  
c) Zero  
d) None
34. According to Bohr's theory the outer orbit electron has \_\_\_\_\_ energy than inner orbits.  
a) Greater  
b) Smaller  
c) Equal  
d) None of these
35. X – rays was discovered by  
a) Bacquerel  
b) Mari – curie  
c) Roentgen  
d) Lane
36. The value of Plank's constant is  
a)  $6.63 \times 10^{-34}$  J.sec  
b)  $6.63 \times 10^{-34}$  J/sec  
c)  $6.63 \times 10^{-34}$  sec/J  
d) None
37. Laser is a device which can produce  
Intense beam of light  
Coherent light  
Monochromatic light  
All
38. When magnetic field is applied in the path X – rays , they will be moving in  
a) Straight line  
b) Circular path  
c) Parabolic path  
d) None
39. The quantized energy of first Bohr orbit of hydrogen atom is  
a) 13.04 eV  
b) 13.6 eV  
c) 13.6 eV  
d) 13.5 eV
40. In LASER principle, a photon produce another photon by the process of  
a) Excitation  
b) De-excite  
c) Ionization  
d) None of these
41. Characteristic X – rays are the X – rays which have  
a) High energy photons  
b) Specific wavelengths  
c) Specific frequencies  
d) All of these
42. In Laser a Meta-stable state is  
a) An excite state  
b) In which an electron is usually stable  
c) In which an electron reside  $10^{-3}$  sec  
d) None of these
43. The Meta-stable state of Helium and Neon is  
a) Different  
b) Identical  
c) Nearly identical  
d) None of these
44. Emission of electrons by metal on heating is called  
a) Secondary emission  
b) Field effect  
c) Photoelectric emission  
d) Thermionic emission
45. The numerical value of ground state energy for H-atom in electron volt is:  
a) -10  
b) 13.6  
c) 10  
d) -13.6

Key Chapter # 20

Q. No.	Ans.	Q. No.	Ans.	Q. No.	Ans.
1	a	16	b	31	c
2	a	17	c	32	b
3	b	18	c	33	b
4	d	19	b	34	a
5	a	20	a	35	c
6	b	21	c	36	a
7	c	22	a	37	d
8	a	23	c	38	a
9	a	24	c	39	b & c
10	d	25	c	40	
11	b	26	c	41	d
12	d	27	d	42	c
13	a	28	c	43	c
14	b	29	c	44	d
15	a	30	c	45	d

## CHAPTER 21: NUCLEAR PHYSICS

1. The energy released by fusion of two deuterons into a He nucleus is about
  - a) 24 MeV
  - b) 200 MeV
  - c) 1.02 MeV
  - d) 7.7 MeV
2. Dr. Abdus Salam unified electromagnetic force and \_\_\_\_\_
  - a) Weak nuclear force
  - b) Strong nuclear force
  - c) Magnetic force
  - d) Gravitational force
3. Which of the following have no charge
  - a)  $\alpha$  - rays
  - b)  $\beta$  - rays
  - c)  $\gamma$  - rays
  - d) cathode rays
4. In Wilson cloud chamber, we use:
  - a) Alcohol vapours
  - b) Neon gas
  - c) Bromine gas
  - d) Water vapours
5. A high potential difference of \_\_\_\_\_ is used in GM counter
  - a) 400 volts
  - b) 1000 volts
  - c) 5000 volts
  - d) 4000 volts
6. One Curie is equal to:
  - a)  $3.70 \times 10^{-10} Bq$
  - b)  $3.70 \times 10^{10} Bq$
  - c) 1 Bq
  - d)  $10^3 Bq$
7. The most useful tracer isotope for the treatment of thyroid gland is:
  - a) Cobalt-60
  - b) Carbon-14
  - c) Iodine-131
  - d) Strontium-90
8. The chemical properties of any element depend on its:
  - a) Number of isotopes
  - b) Number of isobars
  - c) Atomic number
  - d) Mass number
9. The number of protons in any atom are always equal to the number of:
  - a) Electrons
  - b) Neutrons
  - c) Positrons
  - d) Mesons
10. Types of quarks are:
  - a) 4
  - b) 6
  - c) 8
  - d) 10
11.  $\beta$  -particles in Wilson cloud chamber have:
  - a) Zigzag or erratic path
  - b) Curved path
  - c) Circular path
  - d) Elliptical path
12. Nuclear fission chain reaction is controlled by using:
  - a) Steel rods
  - b) Graphite rods
  - c) Cadmium rods
  - d) Platinum rods
13. Extremely penetrating particles are
  - a) Neutrons
  - b)  $\alpha$  -particles
  - c)  $\beta$  -particles
  - d)  $\gamma$  -particles
14. The nuclear reaction taking place in sun is:
  - a) Fission
  - b) Fusion
  - c) Chain
  - d) Alpha decay
15. An  $\alpha$  -particle contains
  - a) 1 proton and 1 neutron
  - b) 2 protons and 2 neutrons
  - c) 3 protons and 3 neutrons
  - d) 4 protons and 4 neutrons
16. Which of the following belong to hadrons group:
  - a) Protons
  - b) Electrons
  - c) Muons
  - d) Neutrinos
17. Number of isotopes of Helium is:
  - a) 2
  - b) 3
  - c) 4
  - d) 5

18. One joule of energy absorbed in a body per kilogram is equal to:  
 a) 1 rad  
 b) One rem  
 c) One gray  
 d) One sievert
19. In nucleus of uranium  $U_{92}^{235}$ , the number of neutrons will be \_\_\_\_\_  
 a) 92  
 b) 235  
 c) 143  
 d) Different for different isotopes
20. One a.m.u is equal to \_\_\_\_\_  
 a)  $1.66 \times 10^{-27}$  kg  
 b)  $1.66 \times 10^{-25}$  kg  
 c)  $1.66 \times 10^{-20}$  kg  
 d) All of above
21. According to which one of following law, the density of nucleus is uniform ?  
 a) J.J. Thomson  
 b) Rutherford's Model  
 c) Bohr's Model  
 d) All of above laws
22. For chain reaction to buildup, the size of the radio active target should be \_\_\_\_\_  
 a) 90  
 b) Greater than the critical size  
 c) Less than the critical size  
 d) Equal to critical size
23. After two half lives, the number of decayed nuclei of an element are:  
 a)  $N$   
 b)  $\frac{N}{2}$   
 c)  $\frac{N}{4}$   
 d)  $\frac{3N}{4}$
24. The examples of antimatter are:  
 a) Antiproton  
 b) Antineutron  
 c) Positron  
 d) All of above
25. Neutron and proton are commonly known as \_\_\_\_\_  
 a) Nucleons  
 b) Meson  
 c) Boson  
 d) Quartz
26. Half life of Radium is 1590 years. In how many years shall the earth loss all his radium due to radioactive decay?  
 a)  $1590 \times 10^6$  years  
 b)  $1590 \times 10^{12}$  years  
 c)  $1590 \times 10^{25}$  years  
 d) Never
27. Which one of the following radiation possesses maximum penetrating power?  
 a)  $\alpha$  - rays  
 b)  $\beta$  - rays  
 c)  $\gamma$  - rays  
 d) All have equal penetrating power
28. Energy liberated when one atom of U-235 undergoes fission reaction is \_\_\_\_\_  
 a) 200 Mev  
 b) 40 Mev  
 c) 30 Mev  
 d) 20 Mev
29. Nuclear force exist between  
 a) Proton - proton  
 b) Proton - Neutron  
 c) Neutron - Neutron  
 d) All of the above
30. Tick the correct statement  
 a) Moderator slow down the neutron  
 b) Moderator bring the neutrons to rest  
 c) Moderator absorb the neutron  
 d) Moderator reflect the neutron
31. Radioactive decay obeys which one of the following data?  
 a)  $N = N_0 e^{-\lambda t}$   
 b)  $N = N_0 e^{xt}$   
 c)  $N = N_0 e^{-xt/2}$   
 d)  $N_0 = N(Ie^{xt})$
32. Which one of the following possesses maximum velocity?  
 a)  $\alpha$  - rays  
 b)  $\beta$  - rays  
 c)  $\gamma$  - rays  
 d) All of the above have same speed
33. Charge on an electron was determine by \_\_\_\_\_  
 a) Ampere  
 b) Maxwell  
 c) Milliken  
 d) Thomson
34. Charge on neutron is \_\_\_\_\_  
 a)  $+1.6 \times 10^{-19}c$   
 b)  $-1.6 \times 10^{-19}c$   
 c) Zero  
 d) No definite charge

35. A particle having the mass of an electron and the charge of a proton is called  
 a) Antiproton  
 b) Positron  
 c) Gamma rays  
 d) Photon
36. Mass of neutron is \_\_\_\_\_  
 a)  $1.67 \times 10^{-13}$  Kg  
 b)  $1.67 \times 10^{-27}$  Kg  
 c)  $9.1 \times 10^{-31}$  Kg  
 d)  $1.67 \times 10^{-19}$  Kg
37. Nuclei having the same mass number but different atomic number are \_\_\_\_\_  
 a) Isotopes  
 b) Isobars  
 c) Isotones  
 d) Isomers
38. A mass spectrograph sorts out \_\_\_\_\_  
 a) Molecules  
 b) Ions  
 c) Elements  
 d) Isotopes
39. Sum of the masses of constituent nucleons as compared to the mass of the resultant nucleus is \_\_\_\_\_  
 a) Smaller  
 b) Greater  
 c) Same  
 d) Some times smaller some times greater
40. An  $\alpha$  - particle is emitted from  ${}_{88}\text{Ra}^{226}$ , what is the mass and atomic number of the daughter nucleus?
- |    | Mass Number | Atomic Number |
|----|-------------|---------------|
| a) | 224         | 84            |
| b) | 220         | 80            |
| c) | 222         | 86            |
| d) | 226         | 87            |
41. The unit of Radioactivity "Curie" is equal to \_\_\_\_\_  
 a)  $3.74 \times 10^9$  disintegration per sec  
 b)  $3.70 \times 10^{10}$  disintegration per sec  
 c)  $3.55 \times 10^{10}$  disintegration per sec  
 d)  $3.60 \times 10^{10}$  disintegration per sec
42. In liquid metal fast breeder reactor, the type of uranium used is \_\_\_\_\_  
 a)  ${}_{92}\text{U}^{235}$   
 b)  ${}_{92}\text{U}^{238}$   
 c)  ${}_{92}\text{U}^{234}$   
 d)  ${}_{92}\text{U}^{239}$
43. Radioactive materials can be identified by measuring their \_\_\_\_\_  
 a) Hardness  
 b) Density  
 c) Mass  
 d) Half life
44. If one or more of the neutrons emitted during fission can be used to build up further fission then the reaction is self sustained and is known as \_\_\_\_\_  
 a) Fission reaction  
 b) Fusion reaction  
 c) Chain reaction  
 d) Chemical reaction
45. Pair production takes place in the vicinity of heavy nucleus so that \_\_\_\_\_  
 a) Net energy is conserved  
 b) Net charge is conserved  
 c) Net momentum is conserved  
 d) All of the above
46. During an encounter with an atom  $\alpha$  - particle knocks out \_\_\_\_\_  
 a) Protons  
 b) Electrons  
 c) Neutrons  
 d) Nothing
47. Which one of the following radiations are suitable for the treatment of an infection in the interior body?  
 a)  $\alpha$  - rays  
 b)  $\beta$  - rays  
 c)  $\gamma$  - rays  
 d) X - rays
48. Various types of cancer are treated by \_\_\_\_\_  
 a) Cobalt 60  
 b) Strontium - 90  
 c) Carbon 14  
 d) Nickel - 63
49. Sterilizations of surgical instrument, medical supplies and bandages can be done by exposing them to a beam of \_\_\_\_\_  
 a)  $\alpha$  - rays  
 b)  $\beta$  - rays  
 c)  $\gamma$  - rays  
 d) 'b' & 'c' have equal antiseptic properties
50. Charge on  $\alpha$  - particle is \_\_\_\_\_  
 a) +1  
 b) +2  
 c) -2  
 d) -1

51. B-particle ionizes an atom \_\_\_\_\_  
a) Through direct collision  
b) Through electrostatic attraction  
c) Through electrostatic repulsion  
d) All of above
52. T.V. sets and microwave oven emit \_\_\_\_\_  
a) X - rays  
b)  $\alpha$  - rays  
c)  $\beta$  - rays  
d)  $\gamma$  - rays
53. A  $\beta$  - particle in a single encounter \_\_\_\_\_  
a) Loses a small fraction of its energy  
b) Loses most of its energy  
c) Loses no energy at all  
d) Loses energy at all
54. Strontium -90 is used as \_\_\_\_\_  
a)  $\beta$  - particle source  
b)  $\alpha$  - particle source  
c)  $\gamma$  - particle source  
d) Neutrons source
55. The penetration power of  $\beta$  - particle as compared to  $\alpha$ -particle is \_\_\_\_\_  
a) 10 times more  
b) 100 times more  
c) 100 times less  
d) 10 times less
56. Geiger counter is suitable for \_\_\_\_\_  
a) Fast counting  
b) Extremely fast counting  
c) Slow counting  
d) All situations
57. An  $\alpha$  - particle can produce fluorescence in \_\_\_\_\_  
a) ZnS  
b) Barium Palatino cyanide  
c) Calcium tunzstate  
d) All of above
58. Pair production cannot take place in vacuum as \_\_\_\_\_ is not conserved  
a) Energy  
b) Charge  
c) Mass  
d) Momentum
59. Average distance covered by  $\alpha$  - particle in air before its ionizing power ceases is called its \_\_\_\_\_  
a) Trajectory  
b) Range  
c) Firing level  
d) Limit
60.  $\gamma$  - rays are electromagnetic waves like \_\_\_\_\_  
a) Normal light  
b) Heat waves  
c) Micro waves  
d) X - rays
61.  $\beta$  -particle ionizes an atom \_\_\_\_\_  
a) Due to electrostatic force of attraction  
b) Due to electrostatic force of repulsion  
c) Due to direct collision  
d) Due to gravitational force
62.  $\beta$  -particles possess greater penetration power then that of  $\alpha$ -particle due to its \_\_\_\_\_  
a) Smaller ionization power  
b) Energy is not conserved  
c) Neither greater nor smaller ionization power  
d) Same ionization power
63. Pair production can take places only with \_\_\_\_\_  
a) X-rays  
b)  $\gamma$  - rays  
c) UV-rays  
d) IR-rays
64. A device for producing high velocity nuclei is \_\_\_\_\_  
a) Cloud chamber  
b) Linear acceleration  
c) A mass spectrograph  
d) Wilson cloud
65. Which one of the following will be better shield against  $\gamma$  - rays?  
a) Ordinary water  
b) Heavy water  
c) Lead  
d) Aluminum
66. The maximum safe limit does for persons working in nuclear power station are \_\_\_\_\_  
a) 1 rem per week  
b) 5 rem per week  
c) 4 rem per week  
d) 3 rem per week
67. Radiations are used for the treatment of skin of a patient is \_\_\_\_\_  
a)  $\alpha$  - rays  
b)  $\beta$  - rays  
c) X - rays  
d)  $\gamma$  - rays

- 68.** Strong nuclear force
- Increase with magnitude of increasing charge
  - Decreases with magnitude of increasing charge
  - Is independent of charge
  - None
- 69.** Complete the reaction
- $${}_Z X^A \rightarrow {}_{Z+1} X + {}_{-1} \beta^0 + \dots + Q$$
- Neutrino
  - Antineutrino
  - $\alpha$  - particle
  - None
- 70.** The half of uranium – 238 is
- $1.67 \times 10^8$  years
  - $3.3 \times 10^9$  years
  - $4.5 \times 10^8$  years
  - $4.5 \times 10^9$  years
- 71.** The  $\alpha$  - particle ionizes the particles in its way and adopt the path which is
- Curved
  - Straight
  - Zig – Zag
  - None of these
- 72.** Which of the following is similar to electron:
- $\alpha$  – rays
  - $\beta$  – rays
  - $\gamma$  – rays
  - Photons
- 73.** The rate of decay of a radioactive substance:
- Remains constant with time
  - Increase with time
  - Decrease with time
  - May increase or decrease with time
- 74.**  $\gamma$  - rays are absorbed by a sheet of
- 1 to 5 mm of lead
  - 1 to 10 mm of lead
  - 5 to 10 mm of lead
  - None of these
- 75.** Tracks obtained by  $\beta$  - particles in Wilson Cloud Chamber is
- Strong Continuous
  - Discontinuous, not straight thin
  - Weak and no definite tracks
  - None of these
- 76.** The dead time of Geiger Muller counter is of the order of
- Micro second
  - Milli second
  - More than millisecond
  - None of these
- 77.** The breakage of  ${}_{92}^{235}U$  produces the fragments as
- Kr and Ba
  - Sn and Mo
  - Xe and Sr
  - All of them
- 78.** The fuel / fuels used in the reactor are nowadays
- Plutonium – 239
  - Uranium – 233
  - Uranium – 235
  - All of these
- 79.** The temperature of the core of the reactor rises to about
- 1000°C
  - 1100°C
  - 1200°C
  - 1300°C
- 80.** Plutonium can be fissioned by
- Slow neutron
  - Fast neutron
  - Very slow neutron
  - None of these
- 81.** Ultraviolet radiation cause
- Sun burn
  - Blindness
  - Skin Cancer
  - All of them
- 82.** Neutrons are particularly more damaging to
- Legs
  - Heart
  - Eyes
  - Brain
- 83.** Radio isotopes can be made easily by bombardment with
- Electrons
  - Protons
  - Neutrons
  - None of these
- 84.** Subatomic particles are divided into
- Photons
  - Leptons
  - Hadrons
  - All of these

85. One amu is equal to \_\_\_\_\_  
 a) 931 MeV  
 b) 9.31 MeV  
 c) 93.1 MeV  
 d) 0.931 MeV
86. Cobalt -60 emits  $\gamma$  –rays of energy \_\_\_\_\_  
 a) 117 MeV  
 b) 11.7 MeV  
 c) 1.17 MeV  
 d) 1.17 KeV
87. Which of the following statements is correct?  
 a) Moderators slow down the neutrons  
 b) Moderators bring the neutrons to rest  
 c) Moderators absorbs the neutrons  
 d) Moderators reflect the neutrons
88. The half life of radioactive element is \_\_\_\_\_  
 a)  $T_{1/2} = \frac{0.693}{\lambda}$   
 b)  $T_{1/2} = 1.43 \lambda$   
 c)  $T_{1/2} = 0.693 \lambda$   
 d) None of these
89. Hadrons are the particle included  
 a) Protons  
 b) Neutrons  
 c) Mesons  
 d) All of these
90. Lepton's particles which experience no strong nuclear force are  
 a) Electrons  
 b) Muons  
 c) Neutrinos  
 d) All of these
91. The charges on the quarks are  
 a) One unit  
 b) Half unit  
 c) Fraction  
 d) None of these
92. Meson is made from  
 a) A pair of quarks  
 b) A pair of anti quarks  
 c) A pair of quarks and anti quarks  
 d) None of these
93. Fission nuclear reaction leads to \_\_\_\_\_ stability.  
 a) Lesser  
 b) Greater  
 c) Medium  
 d) None

94. If a radioactive isotope of silver have a half life of about 7.5 days. After 15 days the remaining isotope of its original is  
 a) 25%  
 b) 50%  
 c) 7.5%  
 d) 15%
95. A nuclide  ${}_{86}R^{220}$  decays to a new nuclide by two  $\alpha$  -emissions, the nuclide S is  
 a)  ${}_{84}S^{212}$   
 b)  ${}_{82}S^{212}$   
 c)  ${}_{80}S^{220}$   
 d) None

## Key Chapter # 21

Q. No.	Ans.	Q. No.	Ans.	Q. No.	Ans.
1	a	33	c	65	c
2	a	34	c	66	b
3	c	35	b	67	b
4	a	36	b	68	c
5	a	37	a	69	b
6	b	38	d	70	d
7	c	39	b	71	b
8	c	40	c	72	b
9	a	41	b	73	c
10	b	42	b	74	b
11	a	43	d	75	b
12	c	44	c	76	b
13	d	45	d	77	d
14	b	46	b	78	d
15	b	47	d	79	a
16	a	48	b	80	b
17	a	49	d	81	d
18	c	50	b	82	
19	c	51	c	83	c
20	a	52	a	84	d
21	a	53	a	85	a
22	b	54	a	86	c
23	d	55	b	87	a
24	d	56	c	88	a
25		57	d	89	d
26	d	58	d	90	d
27	c	59	b	91	c
28	a	60	d	92	c
29	d	61	b	93	b
30	a	62	a	94	a
31	a	63	b	95	b
32	c	64	b		